TERM-1 SAMPLE PAPER SOLVED

MATHEMATICS (BASIC)

Time Allowed: 90 Minutes

Maximum Marks: 40

General Instructions: Same instructions as given in the Sample Paper 1.

SECTION - A

16 marks

(Section A consists of 20 questions of 1 mark each. Any 16 questions are to be attempted.)

 What will be one of the zeroes of p(x) = ax² + bx + c if a + c = b?
 (a) 3
 (b) 0

• •		• • •
(c)	1	(d) – 1

2. How many number of solutions are there for the following pair of linear equation?

x - 2y + 4 = 0 and 3x + 4y + 2 = 0 (a) Unique (b) Infinite

- (c) No solution (d) Two solution
- 3. The total number of students in class X are 54, out of which there are 32 girls and rest are boys. The class teacher has to select one class representative. She writes the name of each student on a separate card and put the cards in one bag. She randomly draw one card from the bag. What is the probability that the name written on card is of a girl?

(a)
$$\frac{7}{27}$$
 (b) $\frac{11}{27}$
(c) $\frac{16}{27}$ (d) $\frac{4}{27}$


- 4. If 2x + 3y = 5 and 3x + 2y = 10, then what is the value of x - y?
 - (a) 3 (b) 4
 - (c) 5 (d) 6

- 5. What is value of 'k' if the point (-3, *k*) divides the line-segment joining the points (-5, -4) and (-2, 3) in a certain ratio?
 - (a) -1 (b) 3 (c) 2 (d) $\frac{2}{3}$
- 6. Kavita decorated her home beautifully with lights on Diwali. She had three strings of blinking lights with different light colours. The lights of first string remain off for 3 seconds, the second string for 5 seconds and the third string for 6 seconds.





The time interval after which lights of the three strings will glow again after switching them on at the same time is :

- (a) 10 seconds (b) 30 seconds
- (c) 60 seconds (d) 90 seconds
- For any event E, if P(E) = 1, the E is called a:
 (a) Equally likely event
 - (b) Impossible event
 - (c) Sure event
 - (d) Mutually exclusive event
- 8. From a square of side 8 cm, two quadrants of a circle of radii 1.4 cm are cut from two corners. Another circle of radius 4.2 cm is also cut from the centre as shown in the figure. Find the area of the remaining (shaded) portion of the square. [Take $\pi = \frac{22}{7}$]



9. The graphs of y = p(x) are given in figure below. Which among the following shows that p(x) has no zero?





10. If the points A (1, 2), O (0, 0) and C (*a*, *b*) are collinear, than:

(c) a = b (d) a = 3b

11. Polynomial $f(x) = x^2 - 5x + k$ has zeroes α and β such that $\alpha - \beta = 1$, then find the value of 4k.

(a) 6	(b) 12
(c) 18	(d) 24

12. Consider points A(4, 3) and B(x, 5) on the circle with centre O(2, 3). Then the value of x is:

- (a) 3 (b) 2
- (c) 1 (d) 0
- **13.** What are the coordinates of the point, which divides the join of the points (5, 0) and (0, 4) in the ratio 2 : 3 internally?


(a) (8, -3)	(b)	(6, 5)

(c)	$\left(3, \frac{8}{5}\right)$	(d) $\left(\frac{5}{2}, 2\right)$

- 14. The events which have equal chances of occuring and no one is preferred over the other are called events:
 - (a) Complementary
 - (b) Probable
 - (c) Equally likely
 - (d) Most likely
- **15.** If the graph would be parallel to x-axis, then its number of zeros for the graph would be:
 - (a) 0 (b) 1
 - (c) more than 1 (d) 2
- **16.** Calculate the distance between the points P(0, 6) and Q(0, -2).

(a) 8 units	(b) 10 units
(c) 6 units	(d) 4 units

- 17. The HCF of two numbers is 9 and their LCM is 2016. If one number is 54, then find the other number?
 (a) 386
 (b) 336
 - (c) 428 (d) 328
- **18.** Evaluate for *x*, if AB || CD in the given figure.



19. In what ratio does *x*-axis divides the join of A(2, -3) and B(5, 6)?

(a) 1 : 1	(b) 2 : 1
(c) 1:2	(d) 1:3


20. What name is given to a largest positive integer that divides given two positive integers?(a) Coprime(b) HCF

()	()
(c) LCM	(d) both (a) and (c)

16 marks

(Section B consists of 20 questions of 1 mark each. Any 16 questions are to be attempted.)

21. In a playground, 4 friends are standing at the points A, B, C and D as shown in given figure, to play a game.



The distance AB is:

- (a) $\sqrt{3}$ units (b) $2\sqrt{3}$ units
- (c) 6 units (d) $3\sqrt{2}$ units
- **22.** Evaluate $\alpha\beta$, if α and β are the zeroes of the polynomial $x^2 + 5x + 8$.
 - (a) 4 (b) 5
 - (c) 16 (d) None of these
- **23.** The decimal representation of $\frac{17}{8}$ will be. (a) Terminating
 - (b) Non-terminating
 - (c) Non-terminating and repeating
 - (d) Non-terminating and non-repeating
- 24. In a given fraction, y is subtracted from the numerator and 2 is added to the denominator it becomes $\frac{1}{2}$. If 7 is subtracted from the numerator and 2 from denominator, then it becomes $\frac{1}{3}$. Then, what is the fraction?

(a)	23 27	(b)	1 5
(c)	<u>15</u> 26	(d)	<u>13</u> 27

- **25.** Which one of the following is an irrational number.
 - (a) $\sqrt{4}$ (b) $3\sqrt{8}$
 - (c) $\sqrt{100}$ (d) $-\sqrt{0.64}$
- **26.** Evaluate $\sin^2 \theta \cos^2 \theta$. If $\sqrt{3} \tan \theta = 3 \sin \theta$, $\theta \neq 0$ and θ is an acute angle.

(a) 1 (b)
$$\frac{1}{3}$$

(c) $-\frac{1}{3}$ (d) -1

27. What is the value of 'k' in the quadratic polynomial $kx^2 + 4x + 3k$, if the sum of the zeroes is equal to their product ?

(a)
$$\frac{-4}{3}$$
 (b) $\frac{2}{3}$
(c) $\frac{1}{0}$ (d) - 5


28. Priyanka, a X standard student, has only ₹ 1 and ₹ 2 coins in her piggy bank. While counting she found that total number of coins are 50 and amount of money with her is ₹ 75. Observing that certain question arises into her mind. She denote the number of ₹ 1 coins by x and ₹ 2 coins by y.



What are the number of \mathfrak{F} 1 coins in her piggy bank?


(a) 10	(b) 20
(c) 22	(d) 25

29. What is the area of shaded region in the given figure where diameter AB is 12 cm long and AB is trisected at points P and Q.




- **30.** What is the decimal representation of $\frac{17}{125}$?
 - (a) 0.136 (b) 0.017 (c) 0.125 (d) 0.163

31. Determine the value of x, if two quadrilaterals ABCD and PQRS are similar.



32. In the graph shown below the number of zeroes of the polynomial are :



33. At what point, does the linear equation y - 2x = 1 intersect the *y*-axis?

(a) (0, 1)	(b) $\left(-\frac{1}{2}, 0\right)$
(c) $\left(0, \frac{14}{5}\right)$	(d) (0, -14)

34. School is organising a webinar for the students of class fifth, sixth and seventh regarding the cyber crime awareness. He asked the teacher incharges of these classes, to arrange all the students in the same hall. There are 84 students in fifth class, 63 students in sixth class and 42 students in seventh class.



What is minimum number of rows in which the students can be seated so that students of same class sit in a same row?

(a) 9	(b) 21
(c) 7	(d) 42

- **35.** What is the area of a sector, whose perimeter is 30 cm and radius of circle is 6.4 cm?
 - (a) 55 cm^2 (b) 38 cm^2
 - (c) 42 cm^2 (d) 57 cm^2
 - **36.** What is the value of 'k' such that k > 0 if the difference between the zeroes of $4x^2 8kx$

+ 9 15 4?	
(a) $\frac{1}{4}$	(b) $\frac{3}{2}$
(c) $\frac{5}{2}$	(d) $\frac{1}{2}$


37. Rohit is playing a game of spinning a wheel. It is a game of chance consists of spinning an arrow which comes to rest pointing at one of the numbers 1, 2, 3, 4, 5, 6, 7 and 8 (figure) and these are equally likely outcomes.



What is the probability that the arrow rest on even number?




38. In the cross country race, the runners had to run along a fixed track. A runner is running along a straight path parallel to a given boundary. This could be represented by an equation of a straight line on a graph paper.



The path of the runner represents the graph of which polynomial?

- (a) Linear polynomial
- (b) Quadratic polynomial
- (c) Cubic polynomial
- (d) None of these
- 39.250 lottery ticket were sold, out of which5 had prizes. If Rahul had purchased onelottery ticket, then the probability that hewins the prize is:
 - (a) $\frac{1}{25}$ (b) $\frac{1}{50}$ (c) $\frac{3}{17}$ (d) $\frac{2}{25}$
- 40. As shown in the figure, ABCD is a square of

side 7 cm and A, B, C and D are centres of equal circles touching externally in pairs. The area of the shaded region is:



8 marks


SECTION - C (Section C consists of 10 questions of 1 mark each. Any 8 questions are to be attempted)

Q 41 to Q 45 Based on Case Study-1: Case Study-1:

Google maps cartography team is working on improving the scalability quality of maps when you use the app on your phones to zoom in using 4 fingers. They are using a proprietary tool called 'MapMaker' to figure out scalability factors. A mathematical model is created for a type of object (below cross-section) to test its scalability on maps app.



In the diagram, AC = 8 cm, CE = 4 cm and the area of the triangle BEC is 4.2 sq cm. Another enlargement with centre E, maps $\triangle EBC$ onto $\triangle EFA$. BC = 3.6 cm.



41. An enlargement , with centre A, maps △ABC onto △ADE, then the scale factor of the enlargement is:

(a)	$\frac{2}{1}$	(b) $\frac{3}{2}$
(c)	<u>4</u> 3	(d) $\frac{2}{3}$

42. The area of $\triangle ABC$ is:


(a) 4.2 sq cm	(b) 6.3 sq cm
(c) 8.4 sq cm	(d) 12.6 sq cm

- **43.** The length of AF is:
 - (a) 7.2 cm (b) 8.4 cm
 - (c) 10.2 cm (d) 10.8 cm
- **44.** The area of \triangle EFA is:
 - (a) 8.4 sq cm (b) 16.8 sq cm
 - (c) 25.2 sq cm (d) 37.8 sq cm
- **45.** The area of \triangle BAF is:
 - (a) 8.4 sq cm (b) 16.8 sq cm
 - (c) 25.2 sq cm (d) 37.8 sq cm

Q 46 to Q 50 Based on Case Study-2: Case Study-2:

Izmir Clock Tower is a historic clock tower in Konak Square in the center of Izmir, Turkey. The French architect Raymond Charles Pere designed the Izmir Clock Tower. It was built in 1901 to commemorate the 25th anniversary of Abdulhamid II's accession to the throne. Four fountains with three water taps each are set around the base of the tower in a circular pattern, and the columns are inspired by Moorish designs. The clock tower has become the symbol of Izmir, and it appeared on the back of Turkish 500 lira banknotes from 1983 to 1989.

Let us assume that the height of the tower AB = 14 m, height of tree CD = 5 m and BD - BC = 1 m. As the tower is vertical, $\angle ABC = 90^\circ$. Further, let us denote $\angle CBD$ by 'a' and $\angle BAD$ by 'b'.



46. The value of sin a	7 is:	
(a) $\frac{12}{13}$	(b) $\frac{13}{12}$	
(c) $\frac{13}{5}$	(d) $\frac{5}{13}$	
47. The value of tan <i>b</i> is :		
(a) $\frac{12}{9}$	(b) $\frac{9}{12}$	
(c) $\frac{15}{12}$	(d) $\frac{15}{9}$	
48. The value of sec ²	$a + \operatorname{cosec}^2 b$ is:	
(a) $\frac{255}{144}$	(b) $\frac{197}{72}$	
(c) 1	(d) $\frac{72}{197}$	
49. The value of sin ² <i>a</i> + cos ² <i>a</i> is :		
(a) 0	(b) –1	
(c) 1	(d) $\frac{1}{4}$	
50. The value of cot ² <i>b</i> is:		
(a) $\frac{81}{144}$	(b) $\frac{144}{225}$	
(c) $\frac{81}{225}$	(d) $\frac{225}{44}$	

SAMPLE PAPER - 11

SECTION - A

1. (d) -1

Explanation: We have, $p(x) = ax^2 + bx + c$

Put
$$x = -1$$
, we get

$$p(-1) = a - b + c$$

$$= (a + c) - b$$

$$= b - b = 0$$
 (as $a + c = b$)

- \therefore -1 is a zero of p(x).
- 2. (d) Unique

Here, pair of equations are : x - 2y + 4 = 0

and 3x + 4y + 2 = 0

Then;

$$\frac{a_1}{a_2} = \frac{1}{3}, \frac{b_1}{b_2} = \frac{-2}{4} = \frac{-1}{2}$$

 $\frac{c_1}{c_2} = \frac{4}{2} = \frac{2}{1}$
 $\therefore \qquad \frac{a_1}{a_2} \neq \frac{b_1}{b_2}$

3. (c) $\frac{16}{27}$ Explanation: Total number of students = 54 Number of girls = 32

$$\therefore P (getting a girl's name) = \frac{32}{54} = \frac{16}{27}$$

4. (c) 5

$$\Rightarrow \qquad x = 4 \\ \therefore \qquad y = -1$$

 $\therefore \qquad x - y = 4 + 1 = 5$

5. (d) $\frac{2}{3}$

Explanation: Let point Q(-3, k) divides AB in the ratio of p: 1.

A(-5, -4)
→ -3 =
$$\frac{-2p-5}{p+1}$$

→ -3p-3 = -2p-5 ⇒ p = 2
∴ Ratio is 2 : 1.
Then, $k = \frac{2 \times 3 - 4}{2 + 1} = \frac{2}{3}$

6. (b) 30 seconds

Explanation: To find the time after which the three lights will glow again, we will find their LCM by prime factorization.

$$3 = 3^{1}$$

 $5 = 5^{1}$
 $6 = 2^{1} \times 3^{1}$

Therefore, LCM = $2^1 \times 3^1 \times 5^1 = 30$


So, that lights of the three strings will glow again together after 30 seconds.

7. (c) Sure event

Explanation: The event with probability 1 is called a sure event, which has a surety of occuring.

8. (b) 5.48 cm²

Explanation: Area of shaded portion = Area of square - 2 × Area of quadrant - Area of circle = $8 \times 8 - 2 \times \frac{1}{4} \times \frac{22}{7} \times 1.4 \times 1.4 - \frac{22}{7} \times 4.2 \times 4.2$ = $64 - 11 \times 0.2 \times 1.4 - 22 \times 0.6 \times 4.2$ = 64 - 3.08 - 55.44= 5.48 cm^2



Explanation: The graph does not intersect *x*-axis at any point. So, it has no zero.

10. (*b*) 2*a* = *b*

Explanation: The given points are O (0, 0), A (1, 2) and C (a, b)

Now, 3 points are collinear

$$\therefore \qquad \text{OC} = \sqrt{(0-a)^2 + (0-b)^2} = \sqrt{a^2 + b^2}$$
$$\text{OA} = \sqrt{(0-1)^2 + (0-2)^2} = \sqrt{1+4} = \sqrt{5}$$
$$\text{AC} = \sqrt{(1-a)^2 + (2-b)^2}$$

Since, points are collinear

$$\therefore \quad AO + OC = AC$$

$$\sqrt{5} + \sqrt{a^2 + b^2} = \sqrt{(1-a)^2 + (2-b)^2}$$

On squaring both sides, we get

$$5 + a^{2} + b^{2} + 2\sqrt{5} \times \sqrt{a^{2} + b^{2}} = 1 + a^{2} - 2a + 4 + b^{2} - 4b$$

-b)²

b

$$\Rightarrow \qquad 2\sqrt{5(a^2+b^2)} = -2(a+2b)$$

On again squaring, we get

$$\Rightarrow \qquad 4a^2 + b^2 - 4ab = 0$$
$$\Rightarrow \qquad (2a - b) = 0 \Rightarrow 2a = 0$$

11. (d) 24

Explanation : Given α and β are the zeroes of the given polynomial.

$$\therefore \qquad \alpha + \beta = -\left(-\frac{5}{1}\right) = 5$$

and
$$\alpha\beta = \frac{k}{1} = k$$

Since,
$$\alpha - \beta = 1$$
$$\Rightarrow \qquad (\alpha - \beta)^2 = 1$$
$$\Rightarrow \qquad (\alpha + \beta)^2 - 4\alpha\beta = 1$$
$$\Rightarrow \qquad (5)^2 - 4 \times k = 1$$
$$\Rightarrow \qquad 25 - 4k = 1$$
$$\Rightarrow \qquad 4k = 24$$

12. (b) 2

Explanation: Since, A and B lie on the circle having centre O. $\bigcirc A = OB$

13. (c) $\left(3, \frac{8}{5}\right)$

Explanation: Let P(x, y) be the point which divides the join of A(5, 0) and B(0, 4) in the ratio 2 : 3 internally.

$$\therefore x = \frac{2(0) + 3(5)}{2+3} = 3 \text{ and } y = \frac{2(4) + 3(0)}{2+3} = \frac{8}{5}$$

Hence, the required point is $\left(3, \frac{8}{5}\right)$

14. (c) Equally likely

Explanation: All the events have equal chances of occuring.

15. (a) 0

Explanation: If the graph would be parallel to *x*-axis it would have no zero, as it will not intersect *x*-axis at any point.

16. (a) 8 units

Explanation: Distance PQ = $\sqrt{(0-0)^2 + (-2-6)^2}$ = 8 units

17. (b) 336

Explanation: Let, the other number be *x*. Then, by formula,

$$LCM \times HCF = 54 \times x$$

$$x = \frac{2016 \times 9}{54}$$
$$= 336$$

18. (b) 7

: ABCD forms a trapezium

$$D$$

$$x - 2$$

$$x + 3$$

$$x + 3$$

$$x + 3$$

$$x + 3$$

$$x + 5$$

$$x + 2$$

$$x + 5$$

$$x + 5$$

$$x + 2$$

$$x + 5$$

$$x + 3$$

19. (c) 1 : 2

Explanation: Let the required ratio be k: 1. We know, *y*-coordinate of any point on *x*-axis is zero.

∴ Using section formula, $\frac{6k-3}{k+1} = 0 \Rightarrow 6k-3 = 0 \Rightarrow k = \frac{1}{2}$ ∴ Required ratio = 1 : 2

20. (b) HCF

Explanation: A largest positive integer that divides given two positive integers completely is called HCF.

21. (d) $3\sqrt{2}$ units **Explanation:** Coordinates of A (3, 4) and B(6, 7)

..
$$AB = \sqrt{(6-3)^2 + (7-4)^2}$$

= $\sqrt{3^2 + 3^2} = \sqrt{9+9}$
= $\sqrt{18} = 3\sqrt{2}$

22. (d) None of these **Explanation:** α and β are the zeroes of the polynomial $x^2 + 5x + 8$.

$$\therefore \qquad \alpha\beta = \frac{8}{1} = 8$$

Hence, none of the given options is correct.

23. (a) terminating

Explanation: As
$$\frac{17}{8} = \frac{17}{2^3} = \frac{17}{2^3 \times 5}$$

Since, denominator is of the form $2^m \times 5^n$, where m and n are integers. So given rational number is a terminating decimal.

24. (c)
$$\frac{15}{26}$$

Explanation: Consider the fraction as $\frac{x}{-}$.

A.T.Q.,
$$\frac{x-1}{y+2} = \frac{1}{2}$$

$$\Rightarrow \quad 2x-2 = y+2$$

$$\Rightarrow \quad 2x-y = 4$$

$$\frac{x-7}{y-2} = \frac{1}{3}$$

$$\Rightarrow \quad 3x-21 = y-2$$

$$\Rightarrow \quad 3x-y = 19$$
On applying (i) - (ii), we get
$$x = 15 \text{ and } y = 26$$

$$\therefore \text{ Required fraction} = \frac{15}{26}$$

25. (b) ³√8

26. (b)
$$\frac{1}{3}$$

Explanation:
We have, $\sqrt{3} \tan \theta = 3 \sin \theta$
 $\Rightarrow \sqrt{3} \frac{\sin \theta}{\cos \theta} = 3 \sin \theta$
 $\Rightarrow \frac{\sqrt{3}}{\cos \theta} = 3$
 $\Rightarrow \frac{\sqrt{3}}{3} = \cos \theta$
 $\therefore \qquad \cos \theta = \frac{1}{\sqrt{3}} \qquad ...(1)$

Now,
$$\sin^2 \theta - \cos^2 \theta = 1 - \cos^2 \theta - \cos^2 \theta$$

= $1 - 2 \cos^2 \theta$
= $1 - 2 \times \left(\frac{1}{\sqrt{3}}\right)^2$ [Using (i)]
= $1 - \frac{2}{3} = \frac{1}{3}$
27. (a) $\frac{-4}{3}$

Explanation: Let α and β be the zeroes of polynomial $kx^2 + 4x + 3k$.

According to the question.

$$\alpha + \beta = \alpha\beta$$
$$\frac{-4}{k} = \frac{3k}{k}$$
$$k = \frac{-4}{3}$$

28. (d) 25

 \Rightarrow

 \Rightarrow

Explanation: The system of linear equations, representing the given situation, is

$$x + y = 50$$
 ...(i)

and
$$x + 2y = 75$$
 ...(ii)

On subtracting (i) from (ii), we get

On substituting y = 25 in (i), we get

$$x = 25$$

Thus, total number of ₹ 1 coins is 25.

Explanation: Here,
$$AP = PQ = QB = \frac{AB}{3} = 4 \text{ cm}$$

 \therefore Area of shaded region = $2 \times \left[\frac{\pi}{2}(4)^2 - \frac{\pi}{2} \times (2)^2\right]$
 $= 2 \times [8\pi - 2\pi]$
 $= 12\pi \text{ cm}^2$

30. (a) 0.136

Explanation: We have,

$$\frac{17}{125} = \frac{17}{5^3} = \frac{17 \times 2^3}{2^3 \times 5^3}$$
$$= \frac{136}{(10)^3} = 0.136$$

31. (b) 6

Explanation : Quadrilaterals ABCD and PQRS are similar.

$$\therefore \qquad \frac{PS}{AD} = \frac{PQ}{AB}$$
(Ratio of corresponding sides are similar)
$$\Rightarrow \qquad \frac{x}{15} = \frac{8}{20}$$

$$\Rightarrow \qquad x = \frac{8 \times 15}{20} = 6$$

32. (d) 3

Explanation: Here, graph touches or cuts the *x*-axis at 3 points. So, number of zeroes are 3.

33. (a) (0, 1)

Explanation: The point at which line cuts the *y*-axis, *x*-coordinate is zero.

 $\therefore \qquad y - 2 \times 0 = 1 \\ y = 1$

.:. Point is (0, 1)

34. (a) 9

Explanation: Here, numbers are 84, 63, and 42 So we need to find their HCF.

 $\therefore \qquad 84 = 2^2 \times 3 \times 7$ $63 = 3^2 \times 7$ $42 = 2 \times 3 \times 7$ $HCF = 3 \times 7$ = 21

:. Number of rows for class fifth = $\frac{84}{21} = 4$

:. Number of rows for class sixth = $\frac{63}{21}$ = 3

:. Number of rows for class seventh = $\frac{42}{21} = 2$

- :. Total number of rows = 4 + 3 + 2 = 9
- **35.** (a) 55 cm²

Explanation: Perimeter of a sector of a circle = 2 × radius + length (*l*) of an arc of sector

 $\therefore \qquad 30 = 2 \times 6.4 + l$ $\Rightarrow \qquad l = 30 - 12.8 = 17.2 \text{ cm}$ Now, area of sector

$$= \frac{1}{2}lr$$
$$= \frac{1}{2} \times 17.2 \times 6.4$$
$$= 55.04 \text{ cm}^2$$
$$\cong 55 \text{ cm}^2 \text{ (approx)}$$

36. (a) $\frac{5}{2}$

Explanation: Here, equation is $4x^2 - 8kx + 9 = 0$ Let, α and β be the zeroes of the given polynomial.

 $\alpha + \beta = -\frac{(-8k)}{4} = -2k$ Then. $\alpha\beta = \frac{9}{4}$ $(\alpha - \beta)^2 = (\alpha + \beta)^2 - 4\alpha\beta$ Since, $4^{2} = (-2k)^{2} - 4 \times \frac{9}{4}$ 16 = 4k² - 9 \Rightarrow $4k^2 = 25$ \Rightarrow $k^2 = \frac{25}{4}$ \Rightarrow $k = \sqrt{\frac{25}{4}} = \pm \frac{5}{2}$ ⇒ But $k = \frac{5}{2}$ *:*..

37. (d) $\frac{1}{2}$

Explanation: \therefore Total number of numbers on the wheel = 8

Let E = Event of getting an even number.

 \therefore Number of outcomes favourable to E = 4

Probability that arrow comes at number 8

$$= P(E) = \frac{Outcomes favourable to E_1 = 4}{Total number of outcomes} = \frac{4}{8}$$

Explanation: As it is a straight line

39. (b)
$$\frac{1}{50}$$

Explanation: Since, total number of tickets = 250

 \therefore Total number of outcomes = 250

Number of tickets having prize *i.e.* favourable outcomes = 5

:. P(getting a prize) =
$$\frac{5}{250}$$

= $\frac{1}{50}$
40. (a) 10.5 cm²

Explanation: Radius of each quadrant

$$=\frac{\text{Side of square}}{2}=\frac{7}{2}\text{cm}$$

:. Area of shaded region = Area of square ABCD – $4 \times$ Area of quadrant.

$$= 7 \times 7 - 4 \times \frac{1}{4} \times \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2}$$
$$= 49 - \frac{77}{2} = 49 - 38.5$$
$$= 10.5 \text{ cm}^2$$

41. (d)
$$\frac{2}{3}$$

Scale factor = $\frac{AC}{AE}$
= $\frac{AC}{AC + CE} = \frac{8}{8 + 4}$
= $\frac{8}{12} = \frac{2}{3}$

42. (d) 12.6 sq. cm

In ΔABC , using Pythagoras theorem, we have

$$AC^{2} = BC^{2} + AB^{2}$$

$$AB^{2} = 64 - 12.96$$

$$= 51.04$$

$$\Rightarrow AB = 7.15 \approx 7.$$

$$∴ Area of △ABC = \frac{1}{2} \times BC \times AB$$

$$= \frac{1}{2} \times 3.6 \times 7$$

$$= 12.6 \text{ cm}^{2}$$

43. (d) 10.8 cm Since, ΔEBC ~ ΔEAF

$$\frac{EC}{EA} = \frac{BC}{AF}$$

$$\Rightarrow \qquad \frac{4}{12} = \frac{3.6}{AF}$$

$$\Rightarrow \qquad AF = 3.6 \times 3$$

$$= 10.8 \text{ cm}$$

44. (d) 37:8 sq. cm

Since, with centre E enlargement as done to ΔEBC and $\Delta \text{EFA}.$

$$\therefore \qquad \Delta \text{EBC} \sim \Delta \text{EFA}$$

$$\frac{ar(\Delta \text{EBC})}{ar(\Delta \text{EFA})} = \frac{\text{EC}^2}{\text{AE}^2}$$

$$\Rightarrow \qquad \frac{4.2}{ar(\Delta \text{EFA})} = \frac{4^2}{12^2}$$

$$\Rightarrow \qquad \text{ar } (\Delta \text{EFA}) = \frac{12 \times 12}{4 \times 4} \times 4.2$$

$$= 3 \times 3 \times 4.2$$

$$= 37.8 \text{ sq. cm}$$

45. (d) 37.8 sq. cm

ar of (
$$\Delta BAF$$
) = $\frac{1}{2} \times AF \times AB$

$$=\frac{1}{2} \times 10.8 \times 7$$


= 37.8 sq. cm

46. (d) $\frac{5}{13}$

Explanation: To find sin *a*, we will first find BD. It is given that BD – BC = 1 m and CD = 5 m. Therefore, applying Pythagoras theorem in triangle BCD, we get : $BD^2 = BC^2 + CD^2 \Rightarrow BD^2 = (BD - 1)^2 + 5^2$ $\Rightarrow BD^2 = BD^2 - 2BD + 1 + 25$ Solving further, 2BD = 26, or BD = 13 m Therefore, BC = 12 m. In $\triangle BCD$, $\sin a = \frac{\text{Perpendicular}}{\text{Hypotenuse}}$ $= \frac{CD}{BD} = \frac{5}{13}$

47. (a) $\frac{12}{9}$

Explanation: To find tan *b*, we will find AE and DE (drawn parallel to BC)



We construct DE || BC and we get a rectangle.

 $\therefore \qquad AB = BE + AE \text{ or } 14 = 5 + AE$ Therefore, AE = 9 m.and, DE = BC = 12 mTherefore, $\tan b = \frac{\text{Perpendicular}}{\text{Hypotenuse}}$ $\frac{DE}{AE} = \frac{12}{9}$

48. (b)
$$\frac{197}{72}$$

Explanation:

sec
$$a = \frac{\text{Hypotenuse}}{\text{Base}}$$

= $\frac{\text{BD}}{\text{BC}} = \frac{13}{12}$
and cosec $b = \frac{\text{Hypotenuse}}{\text{Perpendicular}}$
= $\frac{AD}{\text{DE}} = \frac{15}{12}$

In \triangle AED, by pythagoras theorem AD² = AE² + DE² = 9² + 12² = 81 + 144 = 225 ⇒ AD = 15 m Therefore, $\sec^2 a + \csc^2 b = \left(\frac{13}{12}\right)^2 + \left(\frac{15}{12}\right)^2$ $=\frac{169+225}{144}=\frac{394}{144}=\frac{197}{72}$

49. (c) 1 Explanation: We know,

$$\sin^2 \theta + \cos^2 \theta = 1$$

 \therefore sin²a + cos²a = 1

50. (a)
$$\frac{81}{144}$$

Explanation: $\cot b = \frac{\text{Base}}{\text{Perpendicular}}$
 $= \frac{\text{AE}}{\text{DE}} = \frac{9}{12}$
Therefore, $\cot^2 b = \left(\frac{9}{12}\right)^2 = \frac{81}{144}$